

# Graph Terminology and Special Types of Graphs

Section 10.2

# Basic Terminology


**Definition 1.** Two vertices  $u, v$  in an undirected graph  $G$  are called *adjacent* (or *neighbors*) in  $G$  if there is an edge  $e$  between  $u$  and  $v$ . Such an edge  $e$  is called *incident with* the vertices  $u$  and  $v$  and  $e$  is said to *connect*  $u$  and  $v$ .

**Definition 2.** The set of all neighbors of a vertex  $v$  of  $G = (V, E)$ , denoted by  $N(v)$ , is called the *neighborhood* of  $v$ . If  $A$  is a subset of  $V$ , we denote by  $N(A)$  the set of all vertices in  $G$  that are adjacent to at least one vertex in  $A$ . So,  $N(A) = \bigcup_{v \in A} N(v)$ .

**Definition 3.** The *degree of a vertex in a undirected graph* is the number of edges incident with it, except that a loop at a vertex contributes two to the degree of that vertex. The degree of the vertex  $v$  is denoted by  $\deg(v)$ .

# Degrees and Neighborhoods of Vertices

**Example:** What are the degrees and neighborhoods of the vertices in the graph  $H$ ?



**Solution:**

$H$ :  $\deg(a) = 4$ ,  $\deg(b) = \deg(e) = 6$ ,  $\deg(c) = 1$ ,  
 $\deg(d) = 5$ .

$N(a) = \{b, d, e\}$ ,  $N(b) = \{a, b, c, d, e\}$ ,  $N(c) = \{b\}$ ,

$N(d) = \{a, b, e\}$ ,  $N(e) = \{a, b, d\}$ .

# Handshake Theorem

**Theorem:** Let  $G = (V, E)$  be an undirected graph with  $m$  edges. Then

$$2m = \sum_{v \in V} \deg(v)$$

**Proof:** Each edge contributes twice to the total degree count of all vertices. Thus, both sides of the equation equal to twice the number of edges.

# Handshaking Theorem

We now give two examples illustrating the usefulness of the handshaking theorem.

**Example:** How many edges are there in a graph with 10 vertices of degree six?

**Solution:** Because the sum of the degrees of the vertices is  $6 \cdot 10 = 60$ , the handshaking theorem tells us that  $2m = 60$ . So the number of edges  $m = 30$ .

**Example:** If a graph has 5 vertices, can each vertex have degree 3?

**Solution:** This is not possible by the handshaking theorem, because the sum of the degrees of the vertices  $3 \cdot 5 = 15$  is odd.

# Degree of Vertices

**Theorem:** An undirected graph has an even number of vertices of odd degree.

**Proof:** Let  $V_1$  be the vertices of even degree and  $V_2$  be the vertices of odd degree in an undirected graph  $G = (V, E)$  with  $m$  edges. Then

even  $\rightarrow$   $2m = \sum_{v \in V} \deg(v) = \sum_{v \in V_1} \deg(v) + \sum_{v \in V_2} \deg(v).$

must be even since  $\deg(v)$  is even for each  $v \in V_1$

This sum must be even because  $2m$  is even and the sum of the degrees of the vertices of even degrees is also even. Because this is the sum of the degrees of all vertices of odd degree in the graph, there must be an even number of such vertices.

# Directed Graphs

Recall the definition of a directed graph.


**Definition:** An *directed graph*  $G = (V, E)$  consists of  $V$ , a nonempty set of *vertices* (or *nodes*), and  $E$ , a set of *directed edges* or *arcs*. Each edge is an ordered pair of vertices. The directed edge  $(u, v)$  is said to start at  $u$  and end at  $v$ .

**Definition:** Let  $(u, v)$  be an edge in  $G$ . Then  $u$  is the *initial vertex* of this edge and is *adjacent to*  $v$  and  $v$  is the *terminal* (or *end*) *vertex* of this edge and is *adjacent from*  $u$ . The initial and terminal vertices of a loop are the same.

# Directed Graphs (*continued*)

**Definition:** The *in-degree* of a vertex  $v$ , denoted  $\deg^-(v)$ , is the number of edges which terminate at  $v$ . The *out-degree* of  $v$ , denoted  $\deg^+(v)$ , is the number of edges with  $v$  as their initial vertex. Note that a loop at a vertex contributes 1 to both the in-degree and the out-degree of the vertex.

**Example:** In the graph  $G$  we have



$$\deg^-(a) = 2, \deg^-(b) = 2, \deg^-(c) = 3, \deg^-(d) = 2, \\ \deg^-(e) = 3, \deg^-(f) = 0.$$

$$\deg^+(a) = 4, \deg^+(b) = 1, \deg^+(c) = 2, \deg^+(d) = 2, \\ \deg^+(e) = 3, \deg^+(f) = 0.$$



# Directed Graphs (*continued*)


**Theorem 3:** Let  $G = (V, E)$  be a graph with directed edges. Then:

$$|E| = \sum_{v \in V} \text{deg}^{-}(v) = \sum_{v \in V} \text{deg}^{+}(v).$$

**Proof:** The first sum counts the number of outgoing edges over all vertices and the second sum counts the number of incoming edges over all vertices. It follows that both sums equal the number of edges in the graph. ◀


# Special Types of Simple Graphs: Complete Graphs

A *complete graph on  $n$  vertices*, denoted by  $K_n$ , is the simple graph that contains exactly one edge between each pair of distinct vertices.




# Special Types of Simple Graphs: Cycles and Wheels

A cycle  $C_n$  for  $n \geq 3$  consists of  $n$  vertices  $v_1, v_2, \dots, v_n$ , and edges  $\{v_1, v_2\}, \{v_2, v_3\}, \dots, \{v_{n-1}, v_n\}, \{v_n, v_1\}$ .




A wheel  $W_n$  is obtained by adding an additional vertex to a cycle  $C_n$  for  $n \geq 3$  and connecting this new vertex to each of the  $n$  vertices in  $C_n$  by new edges.



# Special Types of Simple Graphs: $n$ -Cubes


An  $n$ -dimensional hypercube, or  $n$ -cube,  $Q_n$ , is a graph with  $2^n$  vertices representing all bit strings of length  $n$ , where there is an edge between two vertices that differ in exactly one bit position.



# HyperCubes and Parallel Processing

0000	0001	0011	0010
0100	0101	0111	0110
1100	1101	1111	1110
1000	1001	1011	1010


We can map matrices onto this architecture so that neighborhood relations are preserved.



# Bipartite Graphs

**Definition:** A simple graph  $G$  is bipartite if  $V$  can be partitioned into two disjoint subsets  $V_1$  and  $V_2$  such that every edge connects a vertex in  $V_1$  and a vertex in  $V_2$ . In other words, there are no edges which connect two vertices in  $V_1$  or in  $V_2$ .

It is not hard to show that an equivalent definition of a bipartite graph is a graph where it is possible to color the vertices red or blue so that no two adjacent vertices are the same color.





$H$  is not bipartite since if we color  $a$  red, then the adjacent vertices  $f$  and  $b$  must both be blue.

# Bipartite Graphs (*continued*)

**Example:** Show that  $C_6$  is bipartite.


**Solution:** We can partition the vertex set into  $V_1 = \{v_1, v_3, v_5\}$  and  $V_2 = \{v_2, v_4, v_6\}$  so that every edge of  $C_6$  connects a vertex in  $V_1$  and  $V_2$ .



# Bipartite Graphs (*continued*)

**Example:** Show that  $C_3$  is not bipartite.

**Solution:** If we divide the vertex set of  $C_3$  into two nonempty sets, one of the two must contain two vertices. But in  $C_3$  every vertex is connected to every other vertex. Therefore, the two vertices in the same partition are connected. Hence,  $C_3$  is not bipartite.






# Complete Bipartite Graphs

**Definition:** A complete bipartite graph  $K_{m,n}$  is a graph that has its vertex set partitioned into two subsets  $V_1$  of size  $m$  and  $V_2$  of size  $n$  such that there is an edge from every vertex in  $V_1$  to every vertex in  $V_2$ .


**Example:** We display four complete bipartite graphs here.



# New Graphs from Old

**Definition:** A *subgraph* of a graph  $G = (V, E)$  is a graph  $(W, F)$ , where  $W \subset V$  and  $F \subset E$ . A subgraph  $H$  of  $G$  is a proper subgraph of  $G$  if  $H \neq G$ .


**Example:** Here we show  $K_5$  and one of its subgraphs.



# New Graphs from Old

**Definition:** Let  $G = (V, E)$  be a simple graph. The *subgraph induced* by a subset  $W$  of the vertex set  $V$  is the graph  $(W, F)$ , where the edge set  $F$  contains an edge in  $E$  if and only if both endpoints are in  $W$ .

**Example:** Here we show  $K_5$  and the subgraph induced by  $W = \{a, b, c, e\}$ .



# New Graphs from Old (*continued*)

**Definition:** The *union* of two simple graphs  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  is the simple graph with vertex set  $V_1 \cup V_2$  and edge set  $E_1 \cup E_2$ . The union of  $G_1$  and  $G_2$  is denoted by  $G_1 \cup G_2$ .

**Example:**

